arXiv:1110.2237v2 [math.CO] 20 Apr 2012

Upper Bounds on Sets of Orthogonal Colorings of Graphs

Serge C. Ballif

The Pennsylvania State University, University Park, PA 16802

Abstract

We generalize the notion of orthogonal latin squares to colorings of simple graphs. We define two n-colorings of a graph to be *orthogonal* if whenever two vertices share a color in one coloring they have distinct colors in the other coloring. We show that the usual bounds on the maximum size of a certain set of orthogonal latin structures such as latin squares, row latin squares, equi-n squares, single diagonal latin squares, double diagonal latin squares, or sudoku squares are a special cases of bounds on orthogonal colorings of graphs. We also show that the problem of finding a transversal in a latin square of order n is equivalent to finding an n-clique in a particular graph.

Keywords: orthogonal latin squares, graph coloring
2010 MSC: 05B15, 05C15

---

1. Introduction

A *latin square of order n* is an $n \times n$ array filled with n symbols such that each symbol occurs exactly once in each row and column. It is usually convenient to use the numbers $1, \ldots, n$ as the symbols. Two latin squares of order n are said to be *orthogonal* if, when superimposed, each of the n^2 possible ordered pairs of symbols occur. A set of pairwise orthogonal latin squares is said to be a set of *mutually orthogonal latin squares* (MOLS).

We use the notation $N(n)$ for the largest size of a set of MOLS of order n. It is well known that $N(n) \leq n-1$. When n is a power of a prime one can construct a set of $n-1$ MOLS using finite fields, so when n is a prime power $N(n) = n-1$ (see [10, Ch. 2]). In this paper we introduce a generalization of the function $N(n)$ that gives an upper bound on the cardinality of a set of pairwise orthogonal n-colorings of a graph.

Example 1. A *polyomino* is a (possibly disconnected) subset of cells from a square tiling of the plane. If we label the cells of a polyomino so that no symbol occurs more than once in any row or column, then we get a *latin polyomino*. As an example, below we exhibit five latin polyominoes in two polyomino shapes.











Note that the two latin polyominoes on the left are orthogonal in the sense that when they are superimposed each of the ordered pairs is distinct. Similarly, the three latin polyominoes on the right are mutually orthogonal. In each case we have exceeded the bound of $N(2) = 1$ that occurs in the case where the polyomino shape is a square.

In this paper we consider latin squares as a special case of a proper n-coloring of a simple graph. We generalize orthogonality of latin squares to orthogonal colorings of a graph, a notion first studied in [4]. As a corollary, we obtain bounds on the cardinality of orthogonal sets of several structures including equi-n squares, row or column latin squares, latin squares, single diagonal latin squares, and double diagonal latin squares.

A *graph*, G, is a set of *vertices*, $V(G)$, and a set of *edges*, $E(G)$, where each edge is an unordered pair of distinct elements of $V(G)$. The *order* of G is the number of vertices in $V(G)$. Two vertices are said to be *adjacent* (or *neighbors*) if they are connected by an edge in E. We write $u \sim v$ to signify that u is adjacent to V. The *degree* of a vertex is the number of vertices adjacent to it. The *complement* of a graph G is a graph $\overline{G}$ that has vertex set $V(G)$, and two vertices of $\overline{G}$ are adjacent if and only if they are not adjacent in G. An *r-clique* is a (sub)graph of r pairwise adjacent vertices.

A *(proper) coloring* of G is a labeling of the vertices so that any two adjacent edges have distinct labels (which we call *colors*). An *n-coloring* is a coloring consisting of (at most) n colors. It is usually convenient to assume these colors are the numbers $1, \ldots, n$. A graph, G, is said to be *n-colorable* if there exists an n-coloring of G. The *chromatic number* $\chi = \chi(G)$, of G is the minimum number, n, such that there exists an n-coloring of G. If C is a coloring of G, then we write $C(v)$ for the color that C assigns to v.

A latin square is an n-coloring of an $n \times n$ rook's graph. An $m \times n$ *rook's graph* is a graph with mn vertices in m rows and n columns, where two vertices are adjacent if they share a row or a column.

Two colorings, C_1 and C_2, of a graph, G, are said to be *orthogonal* if whenever u and v are distinct vertices of G, we have distinct ordered pairs of colors $(C_1(u), C_2(u)) \neq (C_1(v), C_2(v))$. A set of pairwise orthogonal n-colorings of G is called a set of *mutually orthogonal colorings*. We denote the maximum size of a set of mutually orthogonal n-colorings of a graph G by $N(G, n)$.

Example 2. Below we display orthogonal colorings of a graph of order 9 and a graph of order 16.



A graph with a pair of orthogonal n-colorings can have at most n^2 vertices because there are at most n^2 distinct ordered pairs of labels. In fact, if any color shows up more than n times in a coloring, then the coloring has no orthogonal mate. At the other extreme we may have an overabundance of colors—if $n \geq |V(G)|$ then each vertex can be assigned a different color. Such a coloring is orthogonal to any coloring, including itself.

In the reference, [4], Caro and Yuster investigate the number of colors needed for a graph to have orthogonal colorings. The authors define the *orthogonal chromatic number of G*, $O\chi(G)$, to be the minimum number of colors in any pair of orthogonal colorings of G. Similarly the *k-orthogonal chromatic number of G* is the number, $O\chi_k(G)$, of colors required so that there exist k mutually orthogonal colorings of G. The authors find bounds on the values of $O\chi(G)$ and $O\chi_k(G)$ in terms of the parameters of G such as the maximum degree of any vertex of G or the chromatic number of G. They also show that several classes of graphs always have the lowest possible value $O\chi(G) = \left\lceil \sqrt{|V(G)|} \right\rceil$.

Here we are concerned with the function $N(G,n)$ which is related to the function $O\chi_k(G)$ by the inequalities

$$N(G, O\chi_k(G)) \geq k \qquad \text{and} \qquad O\chi_{N(G,n)}(G) \leq n.$$

In Section 2.4 we obtain some lower bounds for $O\chi_k(G)$ as corollaries to our main theorems.

Many authors have studied edge colorings of graphs as opposed to vertex colorings. A *proper edge coloring* of a graph is a labeling of the edges of a graph such that any two edges that share a vertex receive distinct labels. For each graph, G, the *line graph of G* is the graph with vertex set $E(G)$, and two vertices are adjacent if and only if they share a vertex in G. Each edge coloring of G corresponds to a vertex coloring of the line graph of G in a natural way, so any question about edge coloring can be converted into a question about vertex coloring. The converse statement does not hold because most graphs are not the line graph of any graph.

Despite the fact that each edge coloring may be viewed as an example of vertex coloring, many questions about edge colorings are interesting in their own right and have been studied in the context of edge colorings. Of particular note to us is the article [1] by Archdeacon and Dinitz where they studied orthogonal edge colorings of graphs. In their article they viewed a latin square of order n as an edge coloring of the complete bipartite graph $K_{n,n}$. Results about orthogonal edge colorings can be converted to results about orthogonal vertex coloring by studying the corresponding line graph.

Remark 3. A *subgraph*, H, of a graph, G, is a graph such that $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$. It will always be the case that if H is a subgraph of G, then $N(H,n) \geq N(G,n)$ because mutually orthogonal colorings of G are also mutually orthogonal colorings of H.

Our main results are four theorems that give upper bounds for $N(G,n)$, namely the *clique bound* of Theorem 4, the *degree bound* of Theorem 5, the *average-degree bound* of Theorem 6, and the *edge bound* of Theorem 7. The first two theorems give bounds that depend on a large number of edges locally.

Theorem 4 (clique bound)**.** *Let r, s, and n be integers satisfying $1 < r,s \leq n < r+s$. Let G be a graph with an s-clique, A, that is disjoint from an r-clique, B, such that each vertex in B is adjacent to at least j vertices in A. Then*

$$N(G,n) \leq \left\lfloor \frac{r(s-j)}{r+s-n} \right\rfloor .$$

Theorem 5 (degree bound)**.** *Let G be a graph of order $n^2 - m$ that has l vertices of degree at least d where $0 \leq m \leq l \leq n$. Then*

$$N(G,n) \leq \left\lfloor \frac{(n^2-m-d-1)l}{nl-l-m} \right\rfloor .$$

The next two theorems give upper bounds on $N(G,n)$ based on the number of edges globally.

Theorem 6 (average-degree bound)**.** *Let G be a graph of order v with average vertex degree D. Then for $n < v$,*

$$N(G,n) \leq \left\lfloor \frac{(v-D-1)n}{v-n} \right\rfloor .$$

Theorem 7 (edge bound)**.** *Let G be a graph with v vertices and e edges, $v > n$. Write $v = qn + r$ with $0 \leq r < n$. Then*

$$N(G,n) \leq \left\lfloor \frac{\binom{v}{2} - e}{(n-r)\binom{\lfloor \frac{v}{n} \rfloor}{2} + r\binom{\lceil \frac{v}{n} \rceil}{2}} \right\rfloor .$$

If D is the average degree of the vertices of G, and e is the number of edges in G, then $D = \frac{2e}{V(G)}$, so the edge bound and the average-degree bound depend on the same information. In fact the edge bound will always perform as well as the average-degree bound. We have included the average-degree bound because of its simplicity and because it is sufficiently strong to give tight bounds for most latin structures.

In Section 2 we provide several corollaries to Theorems 4–7. We shall show that the best possible upper bounds on the cardinality of sets of mutually orthogonal latin structures can be derived from the four bounds we have introduced. Table 1

Table 1: We list the upper bounds for $N(G,n)$ (the maximum cardinality of mutually orthogonal sets of latin structures). Upper bounds that are known to be best possible are in bold.

	Degree Bound	Average-Degree Bound	Clique Bound	Edge Bound
Graph G	$\left\lfloor \frac{(n^2-m-d-1)l}{nl-l-m} \right\rfloor$	$\left\lfloor \frac{(v-D-1)n}{v-n} \right\rfloor$	$\left\lfloor \frac{r(s-j)}{r+s-n} \right\rfloor$	$\left\lfloor \frac{\binom{v}{2}-e}{(n-r)\binom{\lfloor \frac{v}{n} \rfloor}{2}+r\binom{\lceil \frac{v}{n} \rceil}{2}} \right\rfloor$
equi-n square	$\mathbf{n+1}$	$\mathbf{n+1}$	—	$\mathbf{n+1}$
row or column latin square of order n	$\mathbf{n}$	$\mathbf{n}$	$\mathbf{n}$	$\mathbf{n}$
latin square of order n	$\mathbf{n-1}$	$\mathbf{n-1}$	$\mathbf{n-1}$	$\mathbf{n-1}$
single diagonal latin square of order n	$\mathbf{n-2}$	$\mathbf{n-2}$	$\mathbf{n-2}$	$\mathbf{n-2}$
double diagonal latin square of odd order, $n > 3$	$\mathbf{n-3}$	$n-2$	$\mathbf{n-3}$	$n-2$
sudoku square of order n	$n-2$	$n-2$	$\mathbf{n-\sqrt{n}}$	$n-2$
m×n latin rectangle $m \leq n$	—	$\mathbf{n-1}$	$\mathbf{n-1}$	$\mathbf{n-1}$

on page 5 summarizes the results of Section 2 by showing the upper bounds on the cardinality of sets of various orthogonal latin structures.

In Section 2.4 we obtain lower bounds on the k-orthogonal chromatic number $O\chi_k(G)$. In Section 3 we prove each of the four bounds. In Section 4 we introduce some graph constructions that arise from sets of mutually orthogonal colorings of a graph.

2. Upper Bounds on sets of Orthogonal Latin Structures

We first define several structures that are related to latin squares. Then we show that the four bounds give the best possible value of $N(G,n)$ for each structure G.

2.1. Latin Squares

An *equi-n square* is an $n \times n$ array such that each of n symbols occurs in exactly n cells. A *row* (resp. *column*) *latin* square of order n is an $n \times n$ array where each of n symbols occurs in each row (resp. column). A *single diagonal latin square* is a latin square where each symbol on the main diagonal is distinct. A *double diagonal latin square* is a single diagonal latin square where each symbol along the back diagonal is also distinct.

Each of these latin structures is a balanced coloring of a graph with n^2 vertices. Two vertices of this underlying graph are adjacent if the structure forbids that they share a color. For instance, two vertices in the underlying graph for a single diagonal latin square are adjacent if they are in the same row or column or if they are both in the main diagonal.

Corollary 8. *Let $n \geq 2$ be a positive integer.*

(1) *No set of mutually orthogonal equi-n squares of order n consists of more than $n+1$ squares.*

(2) *No set of mutually orthogonal row (or column) latin squares of order n consists of more than n squares.*

(3) *No set of mutually orthogonal latin squares consists of more than $n-1$ squares.*

(4) *No set of mutually orthogonal single diagonal latin squares consists of more than $n-2$ squares.*

(5) *No set of mutually orthogonal double diagonal latin squares of odd order $n > 3$ consists of more than $n-3$ squares. (See* [7]*)*

Proof. To prove each part we shall use the degree bound with $l = 1$, $m = 0$, and d equal to the maximum degree of any vertex.

In an equi-n square, no cells are adjacent, so the maximum degree of any vertex in an equi-n square is 0. For statement (1) the degree bound is $n+1$.

In a row latin square the rows form an n-clique, and the maximum degree of any vertex is $n-1$. Thus for (2) the degree bound is $\frac{n^2-1-(n-1)}{n-1} = n$.

In a latin square each vertex is adjacent to all others in the same row and column, so each vertex has degree $2n-2$. Thus for (3) the degree bound is $\frac{n^2-1-(2n-2)}{n-1} = n-1$.

In a single diagonal latin square, the upper left cell is adjacent to each vertex in the top row, the left column, or the main diagonal, so it has degree $3n-3$. In statement (4) we have a degree bound of $\frac{n^2-1-(3n-3)}{n-1} = n-2$.

In a double diagonal latin square of odd order the center cell is adjacent to $4n-4$ cells, so we have a degree bound of $\frac{n^2-1-(4n-4)}{n-1} = n-3$ for (5). ☐

2.2. Gerechte Designs and Sudoku Squares

A *gerechte design of order n* is an $n \times n$ array that has been partitioned into n regions of size n and filled with n symbols so that each row, column, and region contains each symbol exactly once. A *sudoku square of order n^2* is a gerechte design of order n^2 where the partitioned regions are the n^2 subarrays (starting from the top left corner) of size $n \times n$.

As a corollary to Theorem 4 we get a bound on the size of a set of mutually orthogonal gerechte designs. The following bound was proved in [2, Corollary 2.2].

Corollary 9. *Given a partition of an $n \times n$ array into regions $S_1, \ldots, S_n$, each of size n, the size of a set of mutually orthogonal gerechte designs for this partition is at most*

$n-j$, where j is the maximum size of the intersection of a row or column with one of the sets $S_1,\dots,S_n$ with $j<n$.

Proof. Let A be a row (or column). If S_i intersects A in j cells ($j<n$), then we let $B=A\setminus S_i$. Then with $r=|B|$, and $s=t=n$ the clique bound limits the number of mutually orthogonal gerechte designs of order n with partition $S_1,\dots,S_n$ to $\frac{|B|(n-j)}{|B|+n-n}=n-j$. ☐

Corollary 10. *The maximum size of a set of mutually orthogonal sudoku squares of order n^2 is n^2-n.*

Proof. A sudoku square is a gerechte design. The upper left block intersects the top row in n cells, so the clique bound on the cardinality of a set of mutually orthogonal n^2-colorings is precisely n^2-n. ☐

When n is a power of a prime the upper bound of Corollary 10 is always obtainable (see [2] or [11] for constructions). We also note that most of the results of Corollary 8 follow as a corollary to Theorem 4.

2.3. Latin Rectangles

Let $m\le n$. An $m\times n$ *latin rectangle* is defined to be an $m\times n$ array based on n symbols such that no symbol occurs more than once in any row or column. Latin rectangles are a natural extension of the notion of latin squares, and they have been studied extensively [5, 10].

Two $m\times n$ latin rectangles are said to be *orthogonal* if, when superimposed, the mn ordered pairs are distinct. An $m\times n$ latin square is precisely an n-coloring of an $m\times n$ rook's graph, and orthogonality of latin rectangles corresponds precisely with mutually orthogonal n-colorings of those graphs.

It is customary to let $N(m,n)$ be the maximum cardinality of a set of mutually orthogonal latin rectangles (*MOLR*) of size $m\times n$. The following bound for mutually orthogonal latin rectangles is well-known [10].

Corollary 11. $N(m,n)\le n-1$.

Proof. In the underlying graph, G, for an $m\times n$ latin rectangle each vertex shares a row with $n-1$ vertices and column with $m-1$ vertices. Thus we may apply the average-degree bound with $D=m+n-2$ and $v=mn$. Then

$$N(G,n)\le\left\lfloor\frac{(mn-(m+n-2)-1)n}{mn-n}\right\rfloor=n-1.$$ ☐

For $r\le s\le t$ a *latin rectangle of type* (r,s,t) is an $r\times s$ array filled with t distinct symbols such that no symbol occurs more than once in any row or column.

Latin rectangles of type (r,s,t) occur quite naturally. Any $r\times s$ rectangle inside a latin square is a latin rectangle of type (r,s,t) for some t. A latin rectangle of type (n,n,n) is precisely a latin square, and a latin rectangle of type (r,s,s) is an $r\times s$ latin rectangle.

Let $N(r,s,t)$ be the maximum size of a set of MOLR of type (r,s,t). Then $N(r,s) = N(r,s,s)$ because each $r \times s$ latin rectangle is based on s symbols. The bound $N(1,s) = N(1,s,t) = \infty$ is trivial because any two $1 \times s$ latin rectangles are orthogonal. Therefore, we shall assume that $r > 1$.

First we note some obvious bounds. Clearly $N(r,s,t) \geq N(r,s)$ because a rectangle based on s symbols is also based on t symbols because $s \leq t$. In fact with r and s fixed, $N(r,s,t)$ is a nondecreasing function of t. With s and t fixed (and $r < s$) $N(r,s,t)$ is a nonincreasing function of r because a set of mutually orthogonal $(r+1) \times s$ rectangles yields a set of mutually orthogonal $r \times s$ rectangles by deleting a row. Similarly, with r and t fixed (and $r \leq s \leq t$) $N(r,s,t)$ is a nondecreasing function of s because a set of $r \times (s+1)$ MOLR yields a set of $r \times s$ MOLR by deleting a column.

The following result is a direct application of the clique bound of Theorem 4.

Corollary 12. *When $t < 2s$ and $r > 1$,*

$$N(r,s,t) \leq \frac{s^2 - s}{2s - t}.$$

Proof. We use the clique bound of Theorem 4. The top two rows of an $r \times s$ rectangle are s-cliques, A and B. Each vertex in B shares a column with a vertex of A, so we may take $r = s$ and $j = 1$. The clique bound is $\frac{s^2-s}{2s-t}$. $\square$

Note that we get the bound $N(r,s,s) \leq s-1$ of Corollary 11 as a special case of Corollary 12. Also noteworthy is the case when $t = s+1$ where the maximum possible size of a collection of mutually orthogonal latin rectangles of type $(r,s,s+1)$ is s. In particular, when s is one less than a power of a prime the bound is tight because we can use a complete set of s MOLS of order $s+1$ to obtain s mutually orthogonal latin rectangles of type $(r,s,s+1)$ by deleting $r+1$ rows and 1 column from each square.

2.4. Lower Bounds for Orthogonal Chromatic Numbers

In [4] Caro and Yuster studied the the k-orthogonal chromatic number $O\chi_k(G)$, which is the minimum number, n, such that $N(G,n) \geq k$. They obtained upper bounds for $O\chi_k(G)$ that depend on parameters of the graph such as the maximum degree of a vertex or the chromatic number $\chi(G)$. Here we note that we can solve for n in each the four bounds for $N(G,n)$ that we have introduced, and thus find lower bounds for $O\chi_k(G)$.

The following corollary is obtained from the average-degree bound of Theorem 6.

Corollary 13. *Let G be a graph with v vertices with average degree D. Then*

$$O\chi_k(G) \geq \frac{vk}{v - D - 1 - k}$$

The next corollary is obtained from the clique bound of Theorem 4.

Corollary 14. *Let r, s, and n be integers satisfying* $1 < r,s \leq n < r+s$. *Let G be a graph with an s-clique, A, that is disjoint from an r-clique, B, such that each vertex in B is adjacent to at least j vertices in A. Then*

$$O\chi_k(G) \geq \frac{kr+ks+rj-rs}{k}$$

We can also derive lower bounds for $O\chi_k(G)$ from the degree bound and the edge bound, but the formulas are not as clean.

3. Proofs of the Main Theorems

3.1. The Degree Bound

Proof of Theorem 5. Let G be a graph of order n^2-m that has l vertices of degree at least d where $0 \leq m \leq l \leq n$. We must show that

$$N(G,n) \leq \left\lfloor \frac{(n^2-m-d-1)l}{nl-l-m} \right\rfloor .$$

Let $C_1,\ldots,C_k$ be mutually orthogonal n-colorings of G, and let each of the vertices $u_1,\ldots,u_l$ have degree at least d. Since there are n^2-m total vertices, then in any coloring, C_i, there must be at least $n-m$ colors applied to n vertices, while the other m colors are applied to an average of $n-1$ vertices. The number of vertices of one of the colors $C_i(u_1),\ldots,C_i(u_l)$ is at least $m(n-1)+(l-m)n = ln-m$. Thus at least one of the vertices $u_1,\ldots,u_l$ must share a color with $\frac{nl-m}{l}-1$ vertices on average across the colorings $C_1,\ldots,C_k$. Therefore,

$$\left(\frac{nl-m}{l}-1\right)k \leq n^2-m-d-1.$$

Solving for k yields

$$k \leq \frac{(n^2-m-d-1)l}{nl-l-m}. \qquad \square$$

3.2. The Clique Bound

Proof of Theorem 4. Let r, s, and n be integers satisfying $1 < r,s \leq n < r+s$. Let G be a graph with an s-clique, A, that is disjoint from an r-clique, B, such that each vertex in B is adjacent to at least j vertices in A. We must show that

$$N(G,n) \leq \left\lfloor \frac{r(s-j)}{r+s-n} \right\rfloor .$$

Let $C_1,\ldots,C_k$ be mutually orthogonal n-colorings of G. We may assume without loss of generality that the vertices of A are colored with the same colors, say $1,\ldots,s$, so that each pair of colors $(1,1),(2,2),\ldots,(s,s)$ occurs in A for each pair of superimposed colorings C_i, C_j. Among the colorings $C_1,\ldots,C_k$, no vertex of B can have any color in the set $\{1,\ldots,s\}$ more than once, so colors from the set $\{1,\ldots,s\}$

can be applied to a vertex of B at most $s-j$ times across all the colorings $C_1,\dots,C_k$. Thus, colors from the set $\{s+1,\dots,n\}$ will occur at least $k-(s-j)$ times for each of the r vertices of B among the colorings $C_1,\dots,C_k$. The total number of times that colors from the set $\{s+1,\dots,n\}$ occur in B among the colorings $C_1,\dots,C_k$ is at least $r(k-s+j)$.

On the other hand, at most $n-s$ of the colors from the set $\{s+1,\dots,n\}$ can occur in B for each coloring C_i. Thus the total number occurances of a color from the set $\{s+1,\dots,n\}$ in B among the colorings $C_1,\dots,C_k$ is at most $k(n-s)$. Therefore $r(k-s+j)\leq m(n-s)$ so that

$$k \leq \frac{r(s-j)}{r+s-n}. \qquad \square$$

3.3. The Average-Degree Bound

Proof of Theorem 6. Let G be a graph of order v with average vertex degree D. We must show that for $n<v$,

$$N(G,n) \leq \left\lfloor \frac{(v-D-1)n}{v-n} \right\rfloor.$$

Let $C_1,\dots,C_k$ be mutually orthogonal n-colorings of G. First note that if each vertex shares a color with s other vertices in each coloring (on average), then we could argue that $sk+D\leq v-1$ because we can't eliminate every possible pair of vertices from sharing a color since $n<v$. The theorem will be proved by finding a lower bound for s.

Consider a fixed coloring, C_i. We expect that color c will be applied to v/n vertices on average. Therefore, if the color c is selected at random, then we expect each vertex of color c to share a color with $\frac{v}{n}-1$ vertices on average. In other words, if we let l_j be the number of vertices of color j (so $l_1+\cdots+l_n=v$), then the average value of $l_1-1,l_2-1,\dots,l_n-1$ is $\frac{v}{n}-1$.

Next we claim that $s\geq\frac{v}{n}-1$. The actual average number, s, of vertices sharing a color with a given vertex is a weighted average of the values $l_1-1,l_2-1,\dots,l_n-1$. Specifically, the l_i vertices with color i each share a color with l_i-1 vertices, so

$$s=\frac{l_1(l_1-1)+l_2(l_2-1)+\cdots+l_n(l_n-1)}{v}.$$

In this case we have weighted the larger values l_j-1 with greater weights and the smaller values of l_j-1 with lesser weights, so $s\geq\frac{v}{n}-1$. Thus $\left(\frac{v}{n}-1\right)k+D\leq v-1$ so

$$k\leq\frac{(v-D-1)n}{v-n}. \qquad \square$$

3.4. The Edge Bound

Proof of Theorem 7. Let G be a graph with v vertices and e edges, $v>n$. We must show if r is the remainder when v is devided by n, then

$$N(G,n)\leq\left\lfloor\frac{\binom{v}{2}-e}{(n-r)\binom{\lfloor\frac{v}{n}\rfloor}{2}+r\binom{\lceil\frac{v}{n}\rceil}{2}}\right\rfloor.$$

Let $C_1, \ldots, C_k$ be mutually orthogonal n-colorings of G. Each pair of vertices that share a color in C_i must have distinct colors in C_j, $j \neq i$. Thus once each pair of vertices has shared a color in one of the C_i's it is not possible to find an additional n-coloring orthogonal to each of the C_i's. Thus we need to identify the minimum number of pairs of vertices that will share a color in a given C_i. If a color is applied to m vertices, then that accounts for $\binom{m}{2}$ pairs. Therefore, if color t is applied to μ_t vertices, then

$$\binom{\mu_1}{2} + \binom{\mu_2}{2} + \cdots + \binom{\mu_n}{2} \tag{1}$$

pairs of vertices are accounted for in the coloring C_i. The condition $\mu_1 + \cdots + \mu_n = v$ guarantees that the expression (1) is minimized when $n - r$ of the colors appear $\lfloor \frac{v}{n} \rfloor$ times and the other r colors appear $\lceil \frac{v}{n} \rceil$ times. Therefore

$$k \left[(n-r)\binom{\lfloor \frac{v}{n} \rfloor}{2} + r\binom{\lceil \frac{v}{n} \rceil}{2} \right] + e \leq \binom{v}{2}.$$

Solving for k then yields the theorem. □

4. Constructions with Orthogonal Colorings

In this section we introduce a few graph structures that can be created from a set of orthogonal colorings.

4.1. Coloring-Extended Supergraphs

Given a graph G and a collection of colorings $C_1, \ldots, C_k$ of G we can define a graph $G_{C_1,\ldots,C_k}$, which we call the *coloring-extended supergraph* to be the graph with vertex set $V(G)$, and distinct vertices u and v are adjacent in $G_{C_1,\ldots,C_k}$ if either $u \sim v$ in G or $C_i(u) = C_i(v)$ for some i.

Example 15. Below we display two orthogonal colorings, C_1 and C_2, of a graph G and the supergraph G_{C_1,C_2}.



Theorem 16 (Caro and Yuster [4]). *Let G be a graph with mutually orthogonal n-colorings $C_1, \ldots, C_k$. Then the collection, $C_1, \ldots, C_k$, of mutually orthogonal n-colorings of G can be extended if and only if there exists an n-coloring of the graph $G_{C_1,\ldots,C_k}$.*

Proof. Let K be an n-coloring of $G_{C_1,\ldots,C_k}$. Then K is also an n-coloring of G, and $(K(u), C_i(u)) = (K(v), C_i(v))$ implies that $K(u) = K(v)$ and $C_i(u) = C_i(v)$. However, the condition $C_i(u) = C_i(v)$ implies that $u \sim v$ in $G_{C_1,\ldots,C_k}$, so the only possible way to satisfy $K(u) = K(v)$ is if $u = v$. Thus K is orthogonal to each C_i.

For the converse we suppose that K is an n-coloring of G that is orthogonal to each of $C_1, \ldots, C_k$. If $K(u) = K(v)$ then $u \nsim v$ in G, and, since K is orthogonal to C_i, we must have $C_i(u) \neq C_i(v)$ for each i. Therefore K is also an n-coloring of $G_{C_1,\ldots,C_k}$. ☐

Example 17. Below we display orthogonal colorings C and K of a graph G.





Note that K is also a coloring of the graph G_C.

In [4] Caro and Yuster made use of Theorem 16 to find an upper bound for the orthogonal chromatic number $O\chi(G)$.

Remark 18. When $n^2 \geq |G| = v \geq \frac{n^2}{2} + 2$ the clique bound can be applied to the graph G_{C_1}. In this case we obtain the bound

$$N(G,n) \leq \left\lfloor \frac{\left\lceil \frac{v}{n} \right\rceil \left\lceil \frac{v - \left\lceil \frac{v}{n} \right\rceil}{n-1} \right\rceil}{\left\lceil \frac{v}{n} \right\rceil + \left\lceil \frac{v - \left\lceil \frac{v}{n} \right\rceil}{n-1} \right\rceil - n} \right\rfloor + 1.$$

While this is a good bound, it does not outperform the edge bound, though it does match the edge bound for large values of v when the initial graph G has few edges. In particular, when $v = n^2$ we get $N(G,n) \leq n+1$, so part (1) of Corollary 8 can also be viewed as a corollary of the clique bound.

4.2. MOLS via Mutually Orthogonal Colorings

A set of mutually orthogonal colorings of a graph can be used to create a set of MOLS under the right conditions. The existence of $k-2$ mutually orthogonal latin squares of order n is equivalent to the existence of k mutually orthogonal squares of order n. A set of $k-2$ mutually orthogonal latin squares of order n can be extended to a set of k mutually orthogonal equi-n squares by adding a square with constant rows and a square with constant columns. It is similarly possible to construct $k-2$ mutually orthogonal latin squares from a set of k mutually orthogonal equi-n squares. The following theorem and its proof provide a generalization of this construction using mutually orthogonal colorings of graphs.

Theorem 19. *Let G be a graph of order n^2. Suppose there exists a set of k mutually orthogonal n-colorings of G. Then there exists a set of k mutually orthogonal equi-n squares, $k-1$ mutually orthogonal row (or column) latin squares, $k-2$ MOLS of order n, and $k-3$ single diagonal latin squares.*

Proof. We first construct a set of $k-2$ MOLS. Let $C_1, \ldots, C_k$ be mutually orthogonal n-colorings of G. We can obtain a set of $k-2$ MOLS $L_1, \ldots, L_{k-2}$ of order n by letting the entry in row i and column j of L_m be the color $C_m(v)$, where v is the unique vertex in G with color $C_{k-1}(v) = i$ and $C_k(v) = j$. To see that the squares $L_1, \ldots, L_{k-2}$ are latin note that each row and column will have distinct entries because C_{k-1} and C_k are orthogonal colorings. Moreover, since $C_1, \ldots, C_{k-2}$ are pairwise orthogonal, the squares $L_1, \ldots, L_{k-2}$ will also be pairwise orthogonal.

To this set of MOLS we can add an equi-n square with constant columns (or constant rows) to obtain a set of $k-1$ row (or column) latin squares. Moreover, if we include both the square with constant rows and the square with constant columns we have a set of k mutually orthogonal equi-n squares.

To obtain a set of $k-3$ mutually orthogonal single diagonal latin squares we can identify a permutation, σ, that will permute the rows of L_{k-2} so that the main diagonal entries of $\sigma(L_{k-2})$ are equal. Then apply σ to each square $L_1, \ldots, L_{k-3}$. The resulting squares $\sigma(L_1), \ldots, \sigma(L_{k-2})$ will still be mutually orthogonal, and each of $\sigma(L_1), \ldots, \sigma(L_{k-3})$ will be a single diagonal latin square because no symbol will occur more than once down the main diagonal. ☐

4.3. Or-Products

There are several ways to define the product of two graphs. The most commonly studied products are the tensor product graph, the cartesian product graph, the strong product graph, and the lexicographical product graph. We refer the reader to the reference [9] for definitions and results about these product graphs. Here we define yet another product graph.

Let G and H be graphs. We define the *or-product* of G and H to be the graph $G \circledcirc H$ that has vertex set $V(G) \times V(H)$ where vertices (u, v) and (u', v') are adjacent in $G \circledcirc H$ if either $u \sim u'$ in G or $v \sim v'$ in H. The or-product is commutative and associative and each of the product graphs named in the paragraph above is a subgraph of the or-product graph. (In fact $G \circledcirc H$ is the complement of the strong product of the complements of G and H.)

Example 20. The Kronecker product of matrices can be used to create a set of MOLS of order mn from a set of MOLS of order m together with a set of MOLS of order n. However, the $mn \times mn$ rook's graph is not the same as the graph that one obtains by taking the graph theoretic cartesian product of an $m \times m$ rook's graph and an $n \times n$ rook's graph (nor is it the same graph as any other product of rooks graphs).

Below we show a 4×4 rook's graph, G, with a vertex, ▨, that is adjacent to each of the vertices of the form ☒ and a 3×3 rooks graph, H, with a vertex, ▧, that is adjacent to each of the vertices of the form ⊞. We have marked the vertices of the graph $G \circledcirc H$ that are adjacent to the vertex ▩ = (▨,▧).



The following result for graphs is a strengthening of the Kronecker product construction of mutually orthogonal latin squares.

Theorem 21. *Let G and H be graphs and m and n be positive integers. If S is a subgraph of $G \circledcirc H$, then*

$$N(S, mn) \geq \min\{N(G, m), N(H, n)\}.$$

Proof. It is sufficient to show that

$$N(G \circledcirc H, mn) \geq \min\{N(G, m), N(H, n)\}.$$

We create mn colors by taking ordered pairs of colors where the first color is a color applied to vertices of G and the second color is a color applied to vertices of H. We may then use a coloring, A, of G and a coloring, B, of H to obtain a coloring AB of $G \circledcirc H$ by assigning to the vertex (u, v) the color $(A(u), B(v))$. To see that AB is a proper coloring of $G \circledcirc H$ we note that if $(u, v) \neq (u', v')$ and $A_iB_i(u, v) = A_iB_i(u', v')$ then $A_i(u) = A_i(u')$ and $B_i(v) = B_i(v')$, so $u \nsim u'$ and $v \nsim v'$ which implies $(u, v) \nsim (u', v')$.

Now assume that we have mutually orthogonal colorings $A_1, \ldots, A_r$ of G and mutually orthogonal colorings $B_1, \ldots, B_r$ of H. We claim that the product colorings $A_1B_1, \ldots, A_rB_r$ are mutually orthogonal.

Suppose that for $i \neq j$ we have equal pairs of colors

$$\big(A_iB_i(u, v), A_jB_j(u, v)\big) = \big(A_iB_i(u', v'), A_jB_j(u', v')\big).$$

To conclude that A_iB_i is orthogonal to A_jB_j, we must show that $(u, v) = (u', v')$.

Since $A_iB_i(u, v) = A_iB_i(u', v')$ then $A_i(u) = A_i(u')$ and $B_i(v) = B_i(v')$ while $A_jB_j(u, v) = A_jB_j(u', v')$ implies $A_j(u) = A_j(u')$ and $B_j(v) = B_j(v')$. Since A_i is orthogonal to A_j, the equality of the ordered pairs $(A_i(u), A_j(u)) = (A_i(u'), A_j(u')))$ implies $u = u'$. Simililarly $(B_i(v), B_j(v)) = (B_i(v'), B_j(v'))$ implies $v = v'$. Therefore the coloring A_iB_i is orthogonal to the coloring A_jB_j. ☐

As a special case of Theorem 21 we obtain the well-known lower bound on the size of a set of MOLS that can be obtained via a Kronecker product construction [10].

Corollary 22. $N(mn) \geq \min\{N(m), N(n)\}$.

Proof. Let R_j denote the $j \times j$ rook's graph for each j. The graph, R_{mn}, is a subgraph of $R_m \circledcirc R_n$ in a natural way, so by Theorem 21

$$N(R_{mn}, mn) \geq \min\{N(R_m, m), N(R_n, n)\}. \qquad \square$$

5. Transverals and Extremal Graph Theory

A *partial transversal of size* r in a latin square is a set of r cells representing each row, column, and symbol at most once, and a *transversal* in a latin square of order n is an partial transversal of size n.

Two of the most famous open problems involving latin squares are Ryser's Conjecture and Brualdi's Conjecture.

Conjecture 23 (Ryser's Conjecture)**.** *Every latin square of odd order possesses a transversal.*

Conjecture 24 (Brualdi's Conjecture)**.** *Every latin square of order n possesses a partial transversal of size* $n-1$.

The best lower bounds on the size of a partial transversal in a latin square are $n - \sqrt{n}$ for $n \leq 7,730,530$ (obtained independently in [12] and [3]) and $n - 11.0525(\log(n))^2$ otherwise [8]. See [6, Ch. 2] for several results on the size of a partial transversal and for a summary of several conjectures related to Ryser's Conjecture and Brualdi's Conjecture.

We say that a set of r vertices is a *partial transversal of size* r of a coloring of G if the r vertices are pairwise non-adjacent and they have r distinct colors in C. An *independent set* of vertices in a graph is a set of pairwise non-adjacent vertices. The complement of an independent set is a clique.

For a coloring, C, of a graph, G, a partial transversal of size r is equivalent to an independent set in the graph G_C or to an r-clique of $\overline{G_C}$. The problem of finding an n-coloring that is orthogonal to C is equivalent to partitioning C into at most n partial transversals.

Thus we can turn the problem of finding a partial transversal in a latin square into the problem of finding a clique in a graph. Unfortunately, the problem of finding the largest clique in a graph is notoriously difficult.

There is a theorem that gives a lower bound on the size of the largest clique in a graph with a given number of vertices and edges.

Theorem 25 (Turán's Theorem)**.** *Any graph on n vertices with more than* $\frac{n^2(r-2)}{2(r-1)}$ *edges must contain an* r*-clique.*

We may view a latin square as a coloring, C, of a rook's graph, G. To guarantee the existence of an r-clique in a graph of order n^2 we must have $\frac{n^4(r-2)}{2(r-1)}$ edges. The graph $\overline{G_C}$ has precisely $\frac{n^2(n-1)(n-2)}{2}$ edges. The inequality

$$\frac{n^2(n-1)(n-2)}{2} > \frac{n^4(r-2)}{2(r-1)}$$

implies $n^2(n-1)(n-2)(r-11) > n^4(r-2)$. If we solve for r we get $r \leq \frac{n^2+3n-2}{3n-2} \approx \frac{1}{3}n$. An observant reader will note that $\frac{1}{3}n$ is significantly lower than these best known lower bounds, so this approach has its limitations. At this point it is unclear whether Turán's Theorem can be modified to take into account the highly uniform structure of the graph $\overline{G_C}$ so that it might be useful in proving the conjectures of Ryser and Brualdi.

Acknowledgements

The author is indebted to Yu Caro for pointing out several relevant references.

References

[1] D. Archdeacon, J. Dinitz, and F. Harary. Orthogonal edge colorings of graphs. In *Proceedings of the sixteenth Southeastern international conference on combinatorics, graph theory and computing (Boca Raton, Fla., 1985)*, volume 47, pages 49–67, 1985.

[2] R. A. Bailey, P. J. Cameron, and R. Connelly. Sudoku, gerechte designs, resolutions, affine space, spreads, reguli, and Hamming codes. *Amer. Math. Monthly*, 115(5):383–404, 2008.

[3] A. E. Brouwer, A. J. Vries, and R. M. A. Wieringa. A lower bound for the length of partial transversals in a latin square:(preprint). 1978.

[4] Y. Caro and R. Yuster. Orthogonal colorings of graphs. *Electron. J. Combin.*, 6:Research Paper 5, 14 pp. (electronic), 1999.

[5] J. Dénes and A. D. Keedwell. *Latin Squares and Their Applications*. Academic Press, New York, 1974.

[6] J. Dénes and A. D. Keedwell. *Latin Squares*, volume 46 of *Annals of Discrete Mathematics*. North-Holland Publishing Co., Amsterdam, 1991. New developments in the theory and applications, With contributions by G. B. Belyavskaya, A. E. Brouwer, T. Evans, K. Heinrich, C. C. Lindner and D. A. Preece, With a foreword by Paul Erdős.

[7] E. Gergely. A remark on doubly diagonalized orthogonal Latin squares. *Discrete Math.*, 10:185–188, 1974.

[8] P. Hatami and P. W. Shor. A lower bound for the length of a partial transversal in a latin square. *J. Comb. Theory, Ser. A*, 115(7):1103–1113, 2008.

[9] W. Imrich and S. Klavžar. *Product Graphs*. Wiley-Interscience Series in Discrete Mathematics and Optimization. Wiley-Interscience, New York, 2000. Structure and recognition, With a foreword by Peter Winkler.

[10] C. F. Laywine and G. L. Mullen. *Discrete Mathematics Using Latin Squares*. Wiley-Interscience Series in Discrete Mathematics and Optimization. John Wiley & Sons Inc., New York, 1998. A Wiley-Interscience Publication.

[11] R. M. Pedersen and T. L. Vis. Sets of mutually orthogonal Sudoku Latin squares. *College Math. J.*, 40(3):174–180, 2009.

[12] D. E. Woolbright. An $n \times n$ Latin square has a transversal with at least $n - \sqrt{n}$ distinct symbols. *J. Combinatorial Theory Ser. A*, 24(2):235–237, 1978.